

RSOS revisited

G. Takács* and G.M.T. Watts†

Institute for Theoretical Physics, Eötvös University,
Pázmány Péter sétány 1/A, Budapest H-1117

† Department of Mathematics, King's College London
Strand, London WC2R 2LS, United Kingdom

February 8, 2002

Abstract

We investigate the issues of unitarity and reality of the spectrum for the imaginary coupled affine Toda field theories based on $a_1^{(1)}$ and $a_2^{(2)}$ and the perturbed minimal models that arise from their various RSOS restrictions. We show that while all theories based on $a_1^{(1)}$ have real spectra in finite volume, the spectra of $a_2^{(2)}$ models is in general complex, with some exceptions. We also correct the S matrices conjectured earlier for the ϕ_{15} perturbations of minimal models and give evidence for a conjecture that the RSOS spectra can be obtained as suitable projections of the folded ATFTs in finite volume.

hep-th/0203073

Report-no: KCL-MTH-02-04, ITP-Budapest-579

PACS codes: 11.55.Ds, 11.30.Na, 11.10.Kk

Keywords: unitarity, finite size effects, minimal models, integrable perturbations, kinks

*E-mail: takacs@ludens.elte.hu

†E-mail: gmtw@mth.kcl.ac.uk

1 Introduction

Unitarity plays an important role in quantum field theory (QFT). In many application of QFT, the framework of quantum theory requires a positive definite conserved probability, which is guaranteed by the Hermiticity of the Hamiltonian and the unitarity of the S matrix. Hermiticity also guarantees that the spectrum of the Hamiltonian is real, an important condition if the Hamiltonian is to be interpreted as the energy of the physical system.

However, there are applications of the QFT formalism when this is not a necessary physical requirement: non-unitary QFTs (even those with complex spectra) appear to play an increasing role in the investigation of statistical mechanical systems (e.g. disordered systems [1]).

In this paper we intend to study the simplest cases: the finite-size spectra of models related to imaginary coupled Toda theories $a_1^{(1)}$ (sine-Gordon) and $a_2^{(2)}$ (ZMS), with periodic boundary conditions. Our main aim is to establish conditions under which these theories have real spectra, continuing our earlier work started in [2]. Reality of the spectrum could be interesting for several reasons: (i) it affects the large distance asymptotics of correlation functions (in case of a complex spectrum, these asymptotics can show oscillating behaviour), (ii) it could allow a redefinition of the theory that renders it unitary (although, as we discuss, physical requirements may prevent that, and it is not at all clear whether this can be performed while maintaining the interpretation of the given model as a local QFT). Finite size spectra are also interesting in their own right: periodic boundary conditions can also be thought of as realization of finite temperature.

We start by recalling the general issues involved, and the relationships between the three distinct concepts of probability conservation, unitarity and the reality of the spectrum.

For any imaginary coupled affine Toda field theory (ATFT), there are three different classes of models which can be considered, namely the *original (unfolded, unrestricted)*, the *folded* and the *restricted (RSOS)* models. We define these in section 3 and discuss the reality of the spectrum in these three classes for the theories related to $a_1^{(1)}$ (sine-Gordon) and $a_2^{(2)}$ (ZMS) in turn.

In the case of $a_1^{(1)}$, the finite-size spectrum of the RSOS models turns out to be a *subset* of the spectrum of a suitable folded model – i.e. the state space of the RSOS model can be thought of as a projection of the space of the folded model. Using this relation, we can conclude that the spectra of all $a_1^{(1)}$ related models are real. In this paper we also give evidence that a similar relationship exists between the RSOS and folded models based on $a_2^{(2)}$. However, this is not sufficient to establish the reality of these models. We then turn to transfer matrix arguments and numerical analysis to investigate the spectrum.

In all cases in which our studies find no violation of reality, there are several possibilities remaining:

1. Apart from cases in which the S -matrix can be shown to be Hermitian analytic (e.g. perturbations of unitary minimal models), we only have numerical (and sometimes matrix perturbation theory) results. Since these numerical tests require ‘scanning’ over the full range of rapidity of each particle in a multi-particle state, there is always a chance that such methods miss some region of rapidities in which reality is violated. In addition, numerical diagonalisation always introduces some numerical errors, and so some threshold condition must be defined to separate real cases from nonreal cases. It remains a question whether reality violations below the threshold are genuine or not.
2. We could only examine transfer matrices containing up to five particles, as the numerical calculations become progressively more and more difficult as we increase the number of particles involved. Therefore, it is possible that higher particle transfer matrices would introduce further constraints on the reality of the spectrum.
3. In the case of $a_2^{(2)}$, our methods only allow us to determine the large volume asymptotics of the spectrum which have a power-like decay as a function of the volume. It is possible that further contributions (decaying exponentially with the volume, e.g. those related to vacuum polarisation) would spoil reality.
4. For $a_2^{(2)}$, the bootstrap is closed only in certain regimes of the parameter space and we do not know the S -matrices of all possible particles in the spectrum. It is possible that reality of the spectrum is only violated in sectors which contain such ‘extra’ particles.

As a result, we can only have definite results in the cases when the spectrum is complex: if we find that some multi-particle transfer matrix has non-phase eigenvalues then we can conclude that the Hamiltonian must have complex eigenvalues. Whenever our examination finds that the spectrum is real we cannot the possibility exclude that further study would find a complex spectrum.

However, the theories for which we cannot find any violation of reality show some very striking patterns, and therefore we believe it is possible that several or possibly all theories falling into these patterns have real spectra. Clearly, further understanding is necessary: the most promising approach would be the development of some exact method to describe the finite size spectra (e.g. an extension of the $a_2^{(2)}$ NLIE [3] to describe excited states) and a deeper understanding of the relation between the folded and the RSOS models in finite volume.

As a side result, we also correct some minor mistakes in our previous paper [2] and the ϕ_{15} S -matrices conjectured earlier by one of us in [4]. We summarise our results in section 5.

2 Unitarity and related concepts in QFT

In this section we discuss the relation between the concepts of probability conservation, unitarity and reality of the spectrum in quantum field theory which are often confused. We also recall the definitions of Hermitian analyticity and R-matrix unitarity and their relationship with unitarity of the S -matrix.

2.1 Unitarity, reality and probabilistic interpretation

For all the theories we consider there exists a non-degenerate sesquilinear form (which we call an inner product for short, even if it is not positive definite) on their space of states which is conserved under the time evolution described by the Hamiltonian i.e. the time evolution operator and the S -matrix preserves this inner product (and the Hamiltonian is Hermitian with respect to the conjugation defined by it). For perturbed conformal field theories such an inner product is inherited from the standard inner product used in CFT, even away from the critical point.

In the cases when this inner product is positive definite (and defines a Hilbert space structure on the space of states) this implies the usual Hermiticity/unitarity, with the consequence that the S -matrix has phase eigenvalues and the energy spectrum is real. This inner product makes possible the usual probabilistic interpretation in quantum theory. These theories are called *unitary* QFTs.

On the other hand, in many systems the inner product is indefinite. It is still possible for the spectrum to be real and the S -matrix eigenvalues to be phases in which case we call the theory a *real* QFTs. However, generally when the inner product is indefinite, then the spectrum is complex and the eigenvalues of the S -matrix are not phases, in which case we call the QFT *nonreal*¹.

One can see that the three notions of unitarity, reality and probability conservation are in fact different. If we want the theory to have a *positive definite conserved probability*, then unitarity is necessary, which is a stronger notion than reality. However, often this is not required; moreover, there exist exotic probability theories which allow for negative or complex probabilities (which in that case of course loses the interpretation in terms of frequency of events).

An important point is whether the S -matrix has any relevance in the case when the theory is non-unitary. It is easier to accept this when the theory is still real, as in the case of the scaling Lee-Yang model (i.e. Virasoro minimal model $(2,5)$ perturbed by its single non-trivial relevant operator ϕ_{13})² or for indeed of any ϕ_{13} perturbations of Virasoro minimal models. However, there are examples of nonreal theories (e.g. the Virasoro minimal model

¹There doesn't seem to be any agreed convention for the naming of what we call real and nonreal theories. We simply use these names in this paper for convenience.

²For an early discussion of this issue, cf. [5].

(3,14) perturbed by operator ϕ_{15} [6]) for which the finite size spectrum extracted using thermodynamic Bethe Ansatz (TBA) for the vacuum energy and the Bethe-Yang equations for multi-particle states matches perfectly the spectrum of the Hamiltonian obtained numerically using truncated conformal space approach (TCSA), even for the complex part of the spectrum.

To summarise, we believe that for a large class of theories (unitary, real and non-real), and in particular for the RSOS models considered in this paper, the spectrum is determined by the S-matrix even when the spectrum is complex.

2.2 Unitarity and Hermitian analyticity

In analytic S -matrix theory the property of unitarity is closely linked with that of Hermitian analyticity [7, 8]. Without entering into details, we intend to recall these concepts and their relations here as they are going to play an important role later.

Unitarity is simply the statement that using a Hermitian conjugation $\dagger$ with respect to a positive definite inner product, the S operator that maps out-states into the in-states has the property

$$SS^\dagger = S^\dagger S = \mathbb{I}. \quad (2.1)$$

Given a theory that is non-unitary but real, since the S operator has phase eigenvalues it is obviously possible to define a new positive definite inner product with respect to which S is unitary. However, this inner product may be inconsistent with the rules of analytic S matrix theory and/or may render the Hamiltonian non-Hermitian (while the Hamiltonian was Hermitian with respect to the original, indefinite inner product). The scaling Lee-Yang model is an example of this situation. Its spectrum contains a single scalar particle which appears as a bound state in the two-particle scattering. S -matrix theory then relates the norm of one-particle states to two-particle states through the residue of the corresponding pole in the two-particle S -matrix. The natural inner product in the perturbed CFT formalism (defined through the Hermiticity of Virasoro generators) is indefinite, with n -particle states having the signature $(-1)^n$ (or $-(-1)^n$, depending on choice, in which case the sign of the residue is consistent with S -matrix theory. However, if one attempts to use a positive definite inner product, then the residue of the two-particle S -matrix at the bound state pole has the “wrong sign”, which is how it is often stated in the literature. As a result, the natural inner product of the scaling Lee-Yang model is indefinite, and it is an example of a *real* but *non-unitary* theory. Therefore the existence of a positive definite redefinition of the inner product does not mean the theory can be made unitary because it may conflict with some other physically motivated requirements. For a further discussion of this issue see [5]. The property that the residues of the bound state poles have the “right sign” is sometimes called “one-particle unitarity” [9].

In integrable theories the whole S operator is encoded in the set of two-particle S -matrices $S_{AB}(\theta)$, where A and B denote the particles (or multiplets if there are internal quantum numbers) and θ is their relative rapidity. It is a simple matter to prove that unitarity of

all possible two-particle S -matrices

$$\sum_{k,l} S_{AB}(\theta)_{ij}^{kl} (S_{AB}(\theta)_{mn}^{kl})^* = \delta_{im} \delta_{jn} , \quad (2.2)$$

(where we explicitly wrote the multiplet indices) implies the unitarity of multi-particle transfer matrices (defined in Appendix C), and also means that the spectrum is real. In writing equation (2.2) we assumed that we had chosen an orthonormal basis in the internal multiplet space.

If, in addition, the inner product on the space of states is positive definite then the theory in question is a unitary QFT. We shall abbreviate the property (2.2) as TU (two-particle unitarity).

Hermitian analyticity (HA) tells us something about the behaviour of the S -matrix elements under complex conjugation. It states that

$$(S_{AB}(\theta)_{ij}^{kl})^* = S_{BA}(-\theta^*)_{lk}^{ji} . \quad (2.3)$$

On the other hand, the S -matrices we investigate here are derived from quantum group R -matrices. R -matrices also satisfy a relation known as “unitarity” in quantum group theory which we call here R -matrix unitarity (RU) and takes the form

$$\sum_{k,l} S_{AB}(\theta)_{ij}^{kl} S_{BA}(-\theta)_{lk}^{nm} = \delta_{im} \delta_{jn} . \quad (2.4)$$

We see that RU and HA together imply TU and thus reality of the spectrum (note that equation (2.2) is meant only for physical i.e. real values of θ). Therefore for the models we consider Hermitian analyticity implies a real spectrum since RU is automatically satisfied due to the quantum group symmetry.

These notions can be appropriately generalised to RSOS theories, where the multi-particle polarisation spaces are not simply tensor products of one-particle ones. Rather, the multiplet structure is labelled by so-called RSOS sequences, which denote ‘the vacua’ between which the particles mediate and are constrained according to so-called ‘adjacency rules’. The two-particle S -matrix therefore carries four vacuum indices a, b, c, d and it is specified in the following way (we omit particle species labels for simplicity):

$$\begin{array}{c} \begin{array}{ccc} & d & \\ \swarrow & & \searrow \\ a & & c \\ \nearrow & & \nwarrow \\ & b & \end{array} \\ \theta \qquad \qquad \phi \end{array} \longleftrightarrow S_{ab}^{cd}(\theta - \phi) . \quad (2.5)$$

For RSOS theories TU, HA and RU take the form:

$$\text{TU : } \sum_d S_{ab}^{cd}(\theta) S_{ae}^{cd}(\theta)^* = \delta_{be} \quad (2.6)$$

$$\text{HA : } S_{ae}^{cd}(\theta)^* = S_{ad}^{ce}(-\theta^*) \quad (2.7)$$

$$\text{RU : } \sum_d S_{ab}^{cd}(\theta) S_{ad}^{ce}(-\theta) = \delta_{be} \quad (2.8)$$

3 The models

The models we study can be classified in the following way:

1. *The original (unfolded, unrestricted) models.* These have an infinite set of degenerate ‘vacua’ and their spectra are built from a fundamental soliton doublet ($a_1^{(1)}$) or triplet ($a_2^{(2)}$) by closing the S -matrix bootstrap.
2. *Folded models.* Using the periodicity of the field theoretic potential, one can choose to identify the ground states after k periods (see [10] for the sine-Gordon case), i.e. in a k -folded model one has k ground states, between which the solitons mediate. The spectrum and the scattering theory for this case can be straightforwardly written down using the well-known S -matrices of the original model.
3. *Restricted (RSOS) models.* At certain ‘rational’ values of the coupling it is possible to define a space of ‘RSOS states’ as a quotient of the full space of states on which the action of the S operator is well defined. These states can be labelled by sequences of ‘vacua’ and the S -matrix factories into 2-particle ‘RSOS type’ S -matrices. These RSOS S -matrices describe the scattering theory of the ϕ_{13} or $\phi_{12}/\phi_{21}/\phi_{15}$ perturbations of Virasoro minimal models, respectively for $a_1^{(1)}$ and $a_2^{(2)}$. These restrictions were discussed in the following papers: the ϕ_{13} case in [11], ϕ_{12}/ϕ_{21} in [12] and ϕ_{15} in [4]. The above construction of the RSOS states relies on the action of the quantum group and has only been defined for the theory on the full line i.e. in infinite spatial volume. However, given the two-particle S -matrix derived in this way one can easily define the theory in finite volume.

All of the models enumerated above are *integrable* and therefore there are numerous results for their finite volume spectra. We shall mainly use results obtained using two methods.

Firstly, the so-called *nonlinear integral equation (NLIE)* approach, which gives exact results for the spectrum and was pioneered by Klümper et al. [13] on the one hand and Destri and de Vega [14] on the other. The original developments concerned mainly the spectra of the $a_1^{(1)}$ case (for the excited states see also [15, 16, 17]).

Secondly, we use the related method of the Bethe-Yang equations (cf. [18] and references therein), which describe the large volume asymptotics of the finite size spectrum. If the NLIE for a given theory is known, then the asymptotics can be derived independently and for $a_1^{(1)}$ it was checked that they agree with the results from the Bethe-Yang equations [16, 17]. In general when the NLIE is not known this is the only known analytic way to obtain information about the finite size spectrum (in some cases, TBA equations are known for the excited states [19], but these cases are covered by the NLIE as well).

In finite volume, since the field theory potential is periodic, it is possible to introduce twisted sectors of the original model [20]. These are labelled by a twist parameter α , defined mod 2π . The twisted sectors have different finite size spectra which are given by a modification of the original NLIE in which α appears explicitly [20].

The finite size spectrum of the k -folded model is the union of the spectra of twisted sectors where the twist runs through the values [10]

$$\alpha = \frac{2\pi m}{k} \quad , \quad m = 0, \dots, k-1 \quad (3.1)$$

When we turn to specific models we shall discuss a relation between the finite size spectra of folded and RSOS models, namely that the finite-size spectrum of an RSOS model is exactly a subset of the spectrum of an appropriate folded model. Such a relation first emerged for the case of $a_1^{(1)}$ [21], and we shall give evidence that a similar relationship holds for $a_2^{(2)}$.

To be precise, for $a_1^{(1)}$ it is known that both the exact spectra (described by the NLIE) and, as a consequence, their asymptotics (described by the Bethe-Yang eqns) of folded and RSOS models are related. For $a_2^{(2)}$ it is known that the exact vacuum energies in finite volume are related (this was shown using the NLIE framework in [3]), but at present the NLIE for excited states is not known. We shall present evidence that the asymptotics of the excited state energies given by the Bethe-Yang eqns are similarly related. We believe similar relations will hold for the exact spectra of every ATFT and the corresponding RSOS models.

There exists some evidence in support of this claim. Al.B. Zamolodchikov has shown that within the perturbed conformal field theory framework, the perturbative expansion for the vacuum energy of the sine-Gordon model with a suitable value of the twist parameter could be reinterpreted as a perturbative expansion in the RSOS model [20], to all orders in perturbation theory. There is also a quantum group argument for the agreement of the vacuum energies [22], which relies on a formulation of the partition function in terms of the system in infinite volume. In addition, using the $a_1^{(1)}$ NLIE it is possible to calculate (a) the exact ultraviolet conformal weights and (b) the energies of the ground state and excited states in the twisted sectors of sine-Gordon model to very high numerical accuracy. Comparison of these data to (a) the known CFT data and (b) numerical finite volume spectra extracted using TCSA, respectively, shows an excellent agreement [21].

It should be possible to find a proof using quantum group arguments directly for the theory in finite volume, by finding an appropriate projection from the folded model to the RSOS space which commutes with the transfer matrices, but we are unaware of such an argument. The knowledge of this projection would also be useful because it could be used to select systematically the RSOS states in the NLIE approach.

We now proceed to give our conventions for the models based on $a_1^{(1)}$ and $a_2^{(2)}$ and give the RSOS—folded relations that we shall check in Section 4.

3.1 Conventions for $a_1^{(1)}$

The $a_1^{(1)}$ theory is simply the sine-Gordon model, for which we take the action to be

$$\mathcal{A}_{sG} = \int d^2x \left(\frac{1}{2} \partial_\nu \Phi \partial^\nu \Phi + \frac{m_0^2}{\beta^2} \cos \beta \Phi \right). \quad (3.2)$$

As is well known, introducing the parameter $q = \exp(8\pi i/\beta^2)$, one can show that the model is invariant with respect to $U_q(a_1^{(1)})$. The spectrum consists of a doublet of solitons and a collection of scalar particles (breathers). As described by [11], the RSOS theory $\mathcal{M}_{p,p'} + \phi_{13}$ is obtained as a restriction at $\beta = \sqrt{8\pi p/p'}$, that is at $q = \exp(\pi i p'/p)$. Putting together the results of [17] and [10], it is straightforward to see that the finite volume spectrum of this RSOS model is a subset of that of the $2p$ -folded model.

3.2 Conventions for $a_2^{(2)}$

We take the action of the $a_2^{(2)}$ model to be

$$\mathcal{A}_{ZMS} = \int d^2x \left[\frac{1}{2} \partial_\nu \Phi \partial^\nu \Phi + \frac{m_0^2}{\gamma} \left(\exp(2i\sqrt{\gamma}\Phi) + 2 \exp(-i\sqrt{\gamma}\Phi) \right) \right]. \quad (3.3)$$

Introducing the parameter

$$q = \exp\left(\frac{i\pi^2}{\gamma}\right) \quad (3.4)$$

one can show that the model has a symmetry under the affine quantum group $\mathcal{U}_q(a_2^{(2)})$ and as a result its S -matrix can be explicitly constructed [12].

The spectrum of the original model consists of a fundamental soliton transforming as a triplet under $U_q(a_1^{(1)})$, having an S -matrix of the form

$$S_{00}^0(x, \gamma) R(x, q), \quad \text{where } x = \exp(2\pi\theta/\xi) \text{ and } \xi = \frac{2\pi}{3} \frac{\gamma}{(2\pi - \gamma)}, \quad (3.5)$$

where S_{00}^0 is a scalar function which is always a pure phase for real x and γ , and $R(x, q)$ is the R -matrix of $U_q(a_2^{(2)})$ in the triplet representation.

Depending on γ , there may be other particles in the spectrum. For $\xi < \pi$ there are breathers B_0^n , $n = 1, \dots, [\pi/\xi]$ formed as bound states of K_0 transforming in the singlet representation, and for $\xi < 2\pi/3$ there are higher kinks K_i , $n = 1, \dots, [2\pi/(3\xi)]$, also formed as bound states of K_0 and transforming in the triplet representation, and their associated breathers B_i^n .

In addition, it is known that there are values of γ for which the bootstrap does not close on this particle content, and for which there are more particles in the spectrum. Since we are only able to investigate the transfer matrices for the kinks K_i and breathers B_i^n , it is only possible to prove that any particular model has a complex spectrum. If we find that the spectrum for these particles is real, that does not imply the reality of the full spectrum simply because the S -matrix bootstrap could still generate S -matrices with non-phase eigenvalues.

The S -matrices of the higher kinks have the form

$$S_{K_m, K_n}(\theta) = S_{m,n}^0(x, \gamma) R((-1)^{m+n}x, q) \quad (3.6)$$

where S_{mn}^0 is a scalar factor.

Since the scalar factors $S_{m,n}^0$ are phases and there are only two different matrix structures for the two-particle S -matrices S_{K_m, K_n} i.e $R(x, q)$ if $m + n$ is even and $R(-x, q)$ if $m + n$ is odd, for the purposes of finding out if the eigenvalues of the transfer matrices (defined in appendix C) it is hence sufficient to consider

- For $\pi/\gamma < 1$: transfer matrices containing only K_0
- For $\pi/\gamma > 1$: transfer matrices containing only K_0 and/or K_1

3.2.1 RSOS models related to $a_2^{(2)}$

The $a_2^{(2)}$ model has two inequivalent RSOS restrictions [12, 4]. From the conjectured ground state NLIE of the $a_2^{(2)}$ -related models [3] one can determine the minimal value of the folding number for which the RSOS ground state is in the spectrum of the folded model. Together with the results in [12, 4], it leads to the following conjectures

1. $\mathcal{M}_{p,p'} + \phi_{12}$ is a projection of the p -folded ZMS model at $\gamma = \pi p/p'$.

The RSOS S -matrices are given in the appendix A where we have specified all the formulae including the $6j$ symbols since the ones we found in the literature all had typos.³

As for the unrestricted case, the matrix part of the S -matrices depend only on q so they depend on p' only modulo $2p$. In fact, we only need study the values $1 \leq p' < p$ since the substitution $p' \rightarrow -p'$ changes the S -matrix elements to their complex

³We are grateful to Giuseppe Mussardo (SISSA) for providing us with a correct set of formulae.

conjugate (if we choose the square root branches in the $6j$ symbols appropriately) and thus simply complex conjugates the transfer matrices and their eigenvalues too as well.

(n.b. the models $\mathcal{M}_{p,p'} + \phi_{21}$ are contained in this class through the identification $\mathcal{M}_{p,p'} + \phi_{21} \equiv \mathcal{M}_{p',p} + \phi_{12}$.)

2. $\mathcal{M}_{p,p'} + \phi_{15}$ is a projection of the 2p-folded ZMS model at $\gamma = 4\pi p/p'$.

The RSOS restriction leading to the S -matrices was performed in [4], however, certain amplitudes had wrong prefactors. The reason was that in the scattering amplitudes of two (oppositely) charged solitons into two neutral ones there is a Clebsh-Gordan factor which was not taken properly into account and therefore the amplitudes of [4] do not satisfy the Yang-Baxter equation and R -matrix unitarity. The corrected amplitudes are listed in Appendix B.

These perturbations lead to renormalisable field theories if the conformal weight of the perturbing field is less than or equal to two, which is equivalent to the condition $\pi/\gamma \geq 1/2$ for both cases, which is also the condition for the unrestricted model to be well-defined.

4 Results

The leading approximation (e.g. in the NLIE formalism) to finite size effects in large volume is given by the Bethe-Yang equations which are summarised in Appendix C. In particular it can be seen that the spectra of the models can only be real if the transfer matrix eigenvalues (denoted in the Appendix by $\lambda^{(s)}(\vartheta|\vartheta_1, \dots, \vartheta_N)$) are all phases for real values of the particle rapidities ϑ_i .

We check for which values of the coupling the various transfer matrices have phase eigenvalues. There are some simplifications we can make.

1. In all our calculation we omit the scalar prefactors of the S -matrices since these are irrelevant for determining whether the eigenvalues are pure phases or not. We call the transfer matrices obtained from the S -matrices without the scalar factor ‘reduced’ transfer matrices.
2. The R -matrices of all models we consider have some discrete symmetries (which we describe here for real values of the rapidity).
 - (a) $R(x, q^*) = R(x, q)^*$, which means that we only need to consider $0 \leq \arg(q) \leq \pi$.
 - (b) $R(x, -1/q) = UR(x, q)U^{-1}$ where U is a diagonal matrix whose nonzero entries are ± 1 .

The reduced transfer matrices are still in general very complicated, even for the original models, and therefore in our study we diagonalised them numerically for a large set of values for the rapidities and other parameters. In all our plots we show only the eigenvalues of the reduced transfer matrices.

4.1 $a_1^{(1)}$ related models

In the case of $a_1^{(1)}$ related models, since the folded and original models are all unitary, their finite size spectra are real and therefore all ϕ_{13} perturbations have real spectra as well, regardless of whether they are perturbations of unitary or non-unitary minimal models. In addition, the NLIE formalism [21] yields manifestly real spectra as well.

In other words:

$$\text{original unitary} \Rightarrow \text{folded unitary} \begin{cases} \Rightarrow \text{RSOS real} \\ \not\Rightarrow \text{RSOS unitary} \end{cases}$$

One might think that, since the RSOS spectrum is simply a subset of the spectrum of the unitary unrestricted model, the RSOS model would simply inherit a positive definite inner product and also be unitary. However, the fact that the spectrum of the RSOS and unrestricted models are different, means that the constraints of analytic S -matrix theory may enforce different inner products. As an example, consider the value $8\pi/\beta^2 = 2/5$. The particle content of the unrestricted model is a soliton doublet $s, \bar{s}$ and a single breather B . The RSOS model, on the contrary, is the Lee-Yang model with a single particle B (the solitons are removed from the spectrum by the RSOS restriction). There is a first order pole in $S_{BB}(\theta)$ at $\theta = 2\pi i/3$ which must have an explanation in terms of on-shell diagrams. In the unrestricted sine-Gordon model, it is explained by the famous Coleman-Thun mechanism [23] as the sum of singular contributions from diagrams with internal soliton lines. In the RSOS model there are no solitons so this pole must be explained by a single diagram in which B occurs as a bound state of two B particles. As we discussed before, the sign of the residue of this pole forces the inner product to be indefinite.

By numerical diagonalisation of the reduced transfer matrices we found that the transfer matrix eigenvalues for the RSOS models $\mathcal{M}_{p,p'} + \phi_{13}$ are a subset of those for the $2p$ -folded sine-Gordon model with $\beta = \sqrt{8\pi p'/p}$. We also observed that the numerically computed transfer matrix eigenvalues are all phases, as expected from the general argument above.

In figure 1, we plot the arguments of the eigenvalues of the reduced two-particle transfer matrices of the RSOS models $(\mathcal{M}_{7,m} + \phi_{13})$ and the 14-folded sine-Gordon model. The RSOS models are only defined for m integer, i.e. for $\arg(q)/\pi$ taking values $m/7$. For these values the eigenvalues of the transfer matrices (shown as blobs) are a subset of the eigenvalues of the matrices of the folded model shown as lines.

The two-particle space of the 14-folded model has dimension 28, and the spectrum consists of 12 eigenvalues each of multiplicity 2 and 4 of multiplicity 1. The RSOS model excludes




Figure 1: Comparison of the eigenvalues of the matrix part of the two-particle transfer matrices for the RSOS model ($\mathcal{M}_{7,m} + \phi_{13}$) and the 14-folded $a_1^{(1)}$ ATFT. The horizontal axis is $\arg(q)/\pi$ restricted to the fundamental domain between 0 and 1, and the vertical axis is the argument of the eigenvalues. The folded models are shown in green/solid lines, and the RSOS models (which are only defined for discrete values of q) are shown as blobs. The relative rapidity of the particles was chosen $\theta = 1/9$.

the eigenvalues of multiplicity 1, 2 of the other eigenvalues, and each of the remaining 10 eigenvalues appears with multiplicity 1 only.

There are RSOS models $\mathcal{M}_{p,p'} + \phi_{13}$ for which the S -matrix is Hermitian analytic (e.g. the unitary case $p' = p + 1$ or the series $p = 2$, see [11]) which then implies that the spectrum is real. Our arguments above, however, extend to *all* ϕ_{13} perturbations.

4.2 $a_2^{(2)}$ related models

In this section, we investigate models related to $a_2^{(2)}$. We start with the 1-folded model where we correct a mistake in [2]. We then consider multi-particle transfer matrices and show the emergence of a pattern which seems to carry over to the folded case and it plays an important role in the RSOS case as well.

For all three classes of model, we must consider separately the case $\pi/\gamma > 1$ for which there are higher kinks, and $\pi/\gamma < 1$ for which there are no higher kinks and consequently fewer constraints on the values of γ allowed by the reality of the spectrum.

We recall that due to the symmetries of the R -matrix, the spectrum of models with $-\pi < \arg(q) < 0$ is obtained by conjugating the spectrum of the model with $1/q$, and so we can restrict our attention to models with $0 \leq \arg(q) \leq \pi$.

4.2.1 The 1-folded models based on $a_2^{(2)}$

We shall consider first the two-particle transfer matrix, and then the higher particle number transfer matrices in turn. As we have mentioned, for our investigation of the reality of the spectrum, we can ignore the scalar prefactors in the S -matrices and need only consider the ‘reduced’ transfer matrices constructed out of the appropriate R -matrices. The two-particle ‘reduced transfer matrices’ are simply the R -matrices themselves: if the kinks being scattered are K_m and K_n , then the appropriate R -matrix is $R((-1)^{m+n}x, q)$.

As calculated in [2], the eigenvalues of $R(x, q)$ are three pairs of doubly degenerate eigenvalues,

$$1, \quad \left(\frac{1 - q^2 \sqrt{x}}{q^2 - \sqrt{x}} \right), \quad \left(\frac{1 + q^2 \sqrt{x}}{q^2 + \sqrt{x}} \right) \quad (4.1)$$

and three eigenvalues λ satisfying

$$\begin{aligned} & (x - q^4)(x + q^6)\lambda^3 + q^6(2 + q^2)(\lambda^2 + x^2\lambda) \\ & + x(q^2 - 1)(1 - 3q^4 + q^8)(\lambda^2 + \lambda) \\ & - q^2(1 + 2q^2)(x^2\lambda^2 + \lambda) + (1 - q^4x)(1 + q^6x) = 0. \end{aligned} \quad (4.2)$$

As stated in [2], for x negative, there are obviously eigenvalues which are not phases, except when $q^4 = 1$. However, in [2] we stated incorrectly that all the eigenvalues were phases for x non-negative and q a phase, and instead the correct result is that there are non-phase eigenvalues for $0 < |\arg(q)/\pi| < 1/4$. This is due to branch cuts that appear in the solutions of the cubic equation (4.2) which we overlooked before.

We were not able to diagonalise any higher particle number transfer matrices exactly. Therefore we resorted to numerical methods for the cases of three, four, five and six particles, combined with matrix perturbation theory in the case of three and four particles.

The 1-folded models with $\pi/\gamma > 1$

If $\pi/\gamma > 1$ then the model contains both K_0 and K_1 . Since the matrix part of S_{K_0, K_1} is $R(-x, q)$, the eigenvalues are just given by (4.1) and (4.2) with x replaced by $-x$, and from looking at (4.1) we see that the two-particle transfer matrix has non-phase eigenvalues unless $q^4 = 1$. This means that, unless $q^4 = 1$, the theory is *nonreal*. We have not investigated the case $q^4 = 1$ any further.

The 1-folded models with $\pi/\gamma < 1$

If $\pi/\gamma < 1$ then the only particles in the spectrum are the fundamental kinks (and the first breather for $\pi/\gamma > 2/3$). In particular, the model contains no higher kinks, and the only reduced transfer matrices to be considered are those constructed out of $R(x, q)$ with x positive.

The eigenvalues of the two-particle case have already been presented, and show that the theory is nonreal for $0 < |\arg(q)/\pi| < 1/4$ and for $3/4 < |\arg(q)/\pi| < 1$.

Numerical investigation of the three-particle transfer matrices show that the theory is also nonreal if $1/4 < |\arg(q)/\pi| < 1/3$ or $2/3 < |\arg(q)/\pi| < 3/4$. It turns out that the strongest violation of reality happens for small x and y , and one can expand the eigenvalues in a perturbation series around $x = y = 0$. Some care must be taken as the $x = y = 0$ transfer matrices have a nontrivial Jordan form, so the Lidskii–Vishik–Lyusternik generalised perturbation theory [24] must be used for the expansion. Without entering into technicalities we only wish to mention that one obtains exactly the same pattern as described above.

Consideration of the four-particle transfer matrix adds further regions of non-reality, $1/3 < |\arg(q)/\pi| < 3/8$ and $5/8 < |\arg(q)/\pi| < 2/3$, and of the five-particle transfer matrix additional regions of $3/8 < |\arg(q)/\pi| < 2/5$ and $3/5 < |\arg(q)/\pi| < 5/8$.

Assuming this pattern to continue for higher particle number transfer matrices, we are led to the conjecture that the eigenvalues of the n particle transfer matrices are always phases for

$$\frac{1}{2} - \frac{1}{2n} \leq \left| \frac{\arg(q)}{\pi} \right| \leq \frac{1}{2} + \frac{1}{2n} \quad (4.3)$$

and at the isolated points

$$\left| \frac{\arg(q)}{\pi} \right| = \frac{1}{2} \pm \frac{1}{2m} \quad , \quad 1 \leq m \leq n. \quad (4.4)$$

and for every other value of q there are non-phase eigenvalues for some value of the rapidities.

This would lead to the result that the spectrum of the 1-folded model is always complex except for the isolated points

$$\left| \frac{\arg(q)}{\pi} \right| = \frac{1}{2} \left(1 \pm \frac{1}{n} \right) \quad (4.5)$$

4.2.2 Higher folded models based on $a_2^{(2)}$

We investigated the folded transfer matrices using numerical diagonalisation and found that the regions of real and complex spectrum are identical in every case to those of the 1-folded models, independent of the folding number.

It is obvious that the regions of complex spectrum contain those of the 1-folded model, since the spectrum of the folded model contains that of the 1-folded model, but it appears that there are no further constraints arising from the twisted sectors.

4.2.3 RSOS models based on $a_2^{(2)}$

The first observation to make is that since the spectrum of the RSOS models is a subset of the spectrum of the appropriate folded model, if the folded model has a real spectrum then so does the RSOS model.

Next, since the RSOS spectrum is a subset of the folded spectrum, it is possible that the RSOS spectrum is real while that of the folded is complex. This is well known to be the case for the perturbations of the unitary minimal models $\mathcal{M}_{p,p\pm 1} + \phi_{12}$ (note that ϕ_{15} is never a relevant perturbation of a unitary minimal model), and we believe that this property is shared by many other models.

As evidence for our assertion that the spectra of the RSOS models are subsets of those of the folded models, in figures 2 and 3 we plot the eigenvalues of the reduced two-particle transfer matrices for K_0 - K_0 scattering of the 10-folded model and of the associated RSOS models (We performed similar checks for the K_0 - K_1 and K_0 - K_0 - K_0 transfer matrices with equally convincing results).

These figures also show that for this particular choice of rapidity difference, there are regions of $\arg(q)/\pi$ for which the folded transfer matrix has non-phase eigenvalues which are included in the regions for which we believe that the folded transfer matrices have non-phase eigenvalues for some values of rapidity. We also see that some of the RSOS restrictions do manage to omit these non-phase eigenvalues, while others do not. We report more fully on our findings in the later sections.

4.2.4 The ϕ_{12} perturbations with $\pi/\gamma > 1$

These theories have at least one higher kink K_1 and we find that the most stringent restrictions already arise from the two-particle transfer matrix involving K_0 and K_1 .

The theories with $p' = \pm 1 \bmod p$ are of the ‘unitary type’: their S -matrices (and transfer matrices) are proportional to those of the unitary models $\mathcal{M}_{p,p+1} + \phi_{12}$ and $\mathcal{M}_{p,p+1} + \phi_{21}$, and since the scattering itself is manifestly unitary the eigenvalues are all phases. Indeed these S -matrices do satisfy Hermitian analyticity which together with R -matrix unitarity implies the S -matrix unitarity equation for the two-particle S -matrices.

Our numerical calculations show that the transfer matrices of the theories $\mathcal{M}_{p,p'} + \phi_{12}$ where $p' = (p \pm 1)/2 \bmod p$ also have phase eigenvalues only, although we do not know any explanation for this fact yet.

Every other theory has non-phase eigenvalues in the K_0 - K_1 transfer matrix; conversely, the models described above still have phase eigenvalues when we consider the three-particle transfer matrices for all combinations of K_0 and K_1 .

We have not tested these models beyond the three-particle transfer matrix. In addition, it is possible that particles arising through the S -matrix bootstrap will also introduce non-phase eigenvalues in the transfer matrices; since the bootstrap has not been completed for




Figure 2: Comparison of the eigenvalues of the matrix part of the two-particle transfer matrices for the RSOS model ($\mathcal{M}_{10,m} + \phi_{12}$) and the 10-folded $a_2^{(2)}$ ATFT. The horizontal axis is $\arg(q)/\pi$ restricted to the fundamental domain between 0 and 1, and the vertical axis is the argument of the eigenvalues. The phase eigenvalues of the folded models are shown in green/thin solid lines, the non-phase eigenvalues in pink/thick solid lines, and the RSOS models (which are only defined for discrete values of q and happen in this case all to be pure phases) are shown as blobs. The relative rapidity of the particles was chosen $\theta = 5$.

the majority of these models, we cannot say anything more about them.

4.2.5 The ϕ_{12} perturbations with $1/2 < \pi/\gamma < 1$

These theories are more usually known as the ‘ ϕ_{21} ’ perturbations. They have no higher kinks in their spectra and at most one breather, and the S -matrix bootstrap closes on these particles. The only possible violation of reality of the spectrum could be introduced by the fundamental kink S -matrix, since the kink-breather and breather-breather S -matrices are pure phases.

We show a selection of our results (up to folding number 50 and particle number 5) in figure 4.

The small (red) points are values which we find definitely to be ‘non-real’. The remaining large (black) points are values which appear to have real spectra up to and including five-particle states. The solid (red, blue and purple) lines are the first three pairs of real series discussed in point 3 below and the dashed (green) lines are models for which a TBA equation for the ground state is known (see below).




Figure 3: Comparison of the eigenvalues of the matrix part of the two-particle transfer matrices for the RSOS model ($\mathcal{M}_{5,m} + \phi_{15}$) and the 10-folded $a_2^{(2)}$ ATFT. The graph is labelled as for figure 2, except that there are now also non-phase eigenvalues of the RSOS model which are shown as black blobs. The relative rapidity of the particles was chosen $\theta = 5$.

We find a complicated pattern of results, with increasingly more models being ruled out by higher and higher particle number transfer matrices. Since we have only investigated transfer matrices involving at most 5 particles, our results are somewhat sketchy but they show the following patterns:

1. All theories for which the folded models have real spectra are obviously real after RSOS restriction, that is those with $|\arg(q)/\pi| = (1 \pm 1/n)/2$. For the range of γ allowed, this gives only the models $\mathcal{M}_{4m,2m+1} + \phi_{12} \equiv \mathcal{M}_{2m+1,4m} + \phi_{21}$ and the models $\mathcal{M}_{2m+1,m+1} + \phi_{12} \equiv \mathcal{M}_{m+1,2m+1} + \phi_{21}$.
2. The perturbations of the unitary minimal models have manifestly unitary S -matrices and real spectra, corresponding to the models $\mathcal{M}_{p+1,p} + \phi_{12} \equiv \mathcal{M}_{p,p+1} + \phi_{21}$.
3. It appears that several other infinite series of models may also have real spectra, at least this appears to be the case from our numerical tests of the transfer matrices up to 5 particles. These models form sequences which tend to the ‘real’ theories described in 1. above. In fact the first example of such a sequence are the unitary models which tend towards the value $\pi/\gamma = 1$, which is the first ‘real’ model with $n = 1$.

The next examples are the series $\mathcal{M}_{4k\pm 1,3k\pm 1} + \phi_{12}$ tending towards $\arg(q)/\pi = 3/4$,




Figure 4: Numerical results for the eigenvalues of the transfer matrices of $\mathcal{M}_{rs} + \phi_{12}$. The heavy (black) points are models for which the multi-particle spectrum appears to be real up to 5 particles, and the light (red) points those which are non-real, with $[\pi/\gamma]=[s/r]$ plotted vertically against r =(folding number). See text for further details

$\mathcal{M}_{3k\pm 1, 2k\pm 1} + \phi_{12}$ tending towards $\arg(q)/\pi = 2/3$, $\mathcal{M}_{8k\pm 3, 5k\pm 2} + \phi_{12}$ tending towards $\arg(q)/\pi = 5/8$, and so on...

These three pairs of series and the unitary models are shown on figure 4 as solid lines.

Some of these models have been considered before in various contexts. The first of these theories, $\mathcal{M}_{3,5} + \phi_{21}$, was considered by G. Mussardo in [25], where he noted that its spectrum was real despite the fact that the S -matrix was ‘non-unitary’.

Since then, (at least) three series of models have been conjectured to have a ground state described by real TBA equations⁴, these being $\mathcal{M}_{m+1, 2m+1} + \phi_{21}$ [26, 27], $\mathcal{M}_{2m+1, 4m} + \phi_{21}$ [28], and $\mathcal{M}_{2m+1, 4m-2} + \phi_{21}$ [29]. These are massive counterparts of the the $|I|=1, 2$ and 4 massless flows considered in [29] with $\pi/\gamma = 1/2 + |I|/(2r)$ and shown in figure 4 as dashed (green) lines. However, as is well known, simply knowing the TBA equations for the ground state does not determine the excited state spectrum uniquely, (a particular example of this effect being the ‘type II’ ambiguity noted in [6]) and the scaling function can describe the ground state of a genuine unitary model and also a ‘non-real’ non-unitary model. So, while TBA equations are a useful alternative way to study the spectrum of a theory, it is not easy to deduce reality properties from the TBA equation for a ground state.

⁴We thank R. Tateo for pointing this out to us.

4.2.6 The ϕ_{15} perturbations with $\pi/\gamma > 1$

These theories again have at least one higher kink K_1 and again we find that the most stringent restrictions already arise from the two-particle transfer matrix involving K_0 and K_1 .

Our numerical analysis shows that only two series of models have the possibility to be real. These are the theories with $p' = 2p \pm 1 \bmod 4p$ for which the S -matrix is Hermitian analytic which guarantees the reality of the spectrum at the level of the Bethe-Yang equations, and the theories with $p' = \pm 1 \bmod 4p$ for which the reality of the spectrum is mysterious.

4.2.7 The ϕ_{15} perturbations with $\pi/\gamma < 1$

Again, these theories have no higher kinks in their spectra, and at most one breather, and the S -matrix bootstrap closes on these particles. The only possible violation of reality of the spectrum could be introduced by the fundamental kink S -matrix, since the kink-breather and breather-breather S -matrices are pure phases.




Figure 5: Numerical results for the eigenvalues of the transfer matrices of $\mathcal{M}_{rs} + \phi_{15}$. The heavy (black) points are models for which the multi-particle spectrum appears to be real up to 4 particles, and the light (red) points those which are non-real, with $\pi/\gamma = s/4r$ plotted vertically against $r = (\text{folding number})/2$. See text for further details

We find an even more complicated pattern of results than for the ϕ_{12} perturbations, once again with increasingly more models being ruled out by higher and higher particle number transfer matrices. In this case the size of the transfer matrices is much larger, and we have only investigated transfer matrices involving at most 4 particles. Our results are even more sketchy but they show the following patterns:

1. All theories for which the folded models have real spectra are still obviously real after RSOS restriction, that is those with $|\arg(q)/\pi| = (1 \pm 1/n)/2$. For the range of γ allowed, this gives only the models $\mathcal{M}_{p,2p+1} + \phi_{15}$ and $\mathcal{M}_{2p+1,4p+4} + \phi_{15}$.
2. Again, it appears that several other infinite series of models may also have real spectra, at least this appears to be the case from our numerical tests of the transfer matrices up to 4 particles. These models form sequences which tend to the ‘real’ theories described in 1. above.

The first example are the theories, $\mathcal{M}_{k,4k-1} + \phi_{15}$ tending towards $\arg(q)/\pi = 1$ and the series $\mathcal{M}_{k,3k\pm 1} + \phi_{15}$ tending towards $\arg(q)/\pi = 3/4$.

At the next level, we find that up to four particles, there are four infinite series tending to $\arg(q)/\pi = 2/3$, namely $\mathcal{M}_{3k\pm 1,8k\pm 3} + \phi_{15}$ and $\mathcal{M}_{6k\pm 1,16k\pm 2} + \phi_{15}$. Since the existence of this second series is in some sense a new phenomenon, we checked that it survives our numerical tests at the five particle transfer matrix level, but we have no opinion whether or not it will survive at all higher particle numbers.

Again, three infinite series of these models have been considered before in various contexts [29]. These series are exactly those which are related to the infinite series of ϕ_{21} perturbations by interchanging the two terms in the ZMS potential as described in e.g. [6]:

$$\begin{aligned}
\mathcal{M}_{2m+1,m+1} + \phi_{12} &\leftrightarrow \mathcal{M}_{2m+1,4m+4} + \phi_{15} , \\
\mathcal{M}_{4m,2m+1} + \phi_{12} &\leftrightarrow \mathcal{M}_{m,2m+1} + \phi_{15} , \\
\mathcal{M}_{4m-2,2m+1} + \phi_{12} &\leftrightarrow \mathcal{M}_{2m-1,4m+2} + \phi_{15} .
\end{aligned} \tag{4.6}$$

These series are shown as dashed (green) lines in figure 5. The final pair of theories is particularly interesting as being an example of a ‘type II’ pair in that these are different theories but share exactly the same ground state scaling function and have a common sub-sector of multi-particle states with identical finite size energy corrections.

5 Conclusions

We have investigated the finite volume spectra of theories based on $a_1^{(1)}$ and $a_2^{(2)}$ imaginary coupled affine Toda field theories. Our main results can be summarised as follows:

1. All theories (original, folded, RSOS) based on $a_1^{(1)}$ have real spectra. To show this, we used the fact that the finite volume spectrum of the RSOS theories can be obtained as suitable projections of folded theories, which are in turn manifestly unitary QFTs related to sine-Gordon theory. We also presented new evidence for this (previously known) correspondence between the folded and RSOS models based on the transfer matrix method.
2. We conjectured that a similar correspondence exists between folded $a_2^{(2)}$ models and RSOS models based on $a_2^{(2)}$, and supported this by numerical diagonalisation of their transfer matrices.
3. We presented substantial evidence that unrestricted (both the folded and the original) $a_2^{(2)}$ models have complex finite volume spectra in general, perhaps with the exception of a few special values of the coupling constant.
4. Similarly, it seems that RSOS theories based on $a_2^{(2)}$ have complex spectra in general, with the exception of some special sequences asymptotically approaching the special values of the coupling for which the unrestricted models may have real spectrum. These sequences in particular include the perturbations of the unitary minimal models, for which we know that the spectrum is real.

There are quite a few open questions remaining. First of all, the NLIE description must be extended to excited states of $a_2^{(2)}$ related models; this could provide us conclusive evidence on whether or not the spectra of the unrestricted $a_2^{(2)}$ theories are real for the special values of the coupling constants for which the transfer matrix calculations found no violation of reality. On the other hand, our results that show that the spectrum is complex in general must be reproduced by such an extension of the NLIE.

Second, the projection of the folded spectrum to the RSOS spectrum must be explicitly realized. Together with the extension of the NLIE to excited states, this would open the way to (a) a systematic description of the spectra of the RSOS models; (b) determining whether the sequences for which we found no violation of reality with our methods really have real spectra.

It seems appropriate to mention that another interesting problem is the relation of the $a_2^{(2)}$ model and ‘ ϕ_{21} ’ perturbations to the corresponding q -state Potts models. See [30] for TBA equations for the ground states of these models and for a discussion of the particle spectra in the q -state Potts models.

Third, as we mentioned in the text, the full spectrum of $a_2^{(2)}$ related theories is not yet closed in full generality. An example is $\mathcal{M}_{3,5} + \phi_{12}$, for which the closed spectrum is not known up to date. One could attempt to close the S matrix bootstrap (we know of some attempts that failed, and we ourselves tried unsuccessfully); but equally well, it could possibly be found by extending the NLIE to describe the excited state spectrum of $a_2^{(2)}$ related theories.

Acknowledgments

The authors would like to thank P. Dorey, H. Saleur and R. Tateo for valuable comments and discussions. We also thank G. Mussardo for providing a correct set of q -6j symbols for the ϕ_{12} perturbations. This work was supported by a Royal Society joint project grant. G.T. is supported by a Zoltán Magyary Fellowship from the Hungarian Ministry of Education, and is partly supported from Hungarian funds OTKA T029802/99 and FKFP-0043/2001. G.T. is also grateful to the Department of Mathematics, King's College, London for their hospitality.

Appendix

A Scattering amplitudes for $\mathcal{M}_{p,p'} + \Phi_{1,2}$

The scattering amplitudes for the fundamental kinks in $\mathcal{M}_{p,p'} + \Phi_{1,2}$ were written down by Smirnov in [12]. Here we briefly recall his result and give explicit expressions for the necessary q -6j symbols as it seems there are none available in the literature that are free of misprints and errors.




Figure 6: Adjacency rules for kinks in Φ_{12} perturbations for $j_{\max} \in \mathbb{Z}$

The allowed vacuum sequences $\{j_1, \dots, j_n\}$ have the following adjacency rules:

$$\begin{aligned} j_{k+1} &= 1 \text{ if } j_k = 0 \\ j_{k+1} &\in \{\max(0, j_k - 1) \dots \min(j_k + 1, p - 3 - j_k)\} \text{ if } j_k \neq 0 \end{aligned} \quad (\text{A.1})$$

and are of two types: either all j_k are integers or all $j_k \in \mathbb{Z} + 1/2$. The maximum value for j_k is $j_{\max} = (p - 2)/2$. The adjacency rules for these two sequences are shown on Figure 6 for the case when $j_{\max}$ is integer; for the case $j_{\max} \in \mathbb{Z} + 1/2$ two very similar graphs result. Let us introduce the notation

$$q = \exp\left(\frac{i\pi p'}{p}\right), \quad x = \exp\left(\frac{2\pi\vartheta}{\xi}\right) \quad \text{where} \quad \xi = \frac{2}{3} \frac{\pi p}{2p' - p}, \quad (\text{A.2})$$

and the q -numbers

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}. \quad (\text{A.3})$$

Then the 2-kink scattering amplitudes take the form

$$\begin{aligned} S_{ab}^{cd}(\vartheta) &= \hat{S}_{ab}^{cd}(x, q) S_0(\vartheta) \\ \hat{S}_{ab}^{cd}(x, q) &= \frac{(q^2 - 1)(q^3 + 1)}{q^{\frac{5}{2}}} \delta_{bd} + \left(\frac{1}{x} - 1\right) q^{\frac{3}{2} - \frac{1}{2}(C_b + C_d - C_a - C_c)} \left\{ \begin{matrix} 1 & c & d \\ 1 & a & b \end{matrix} \right\}_q + \\ &\quad (1 - x) q^{-\frac{3}{2} + \frac{1}{2}(C_b + C_d - C_a - C_c)} \left\{ \begin{matrix} 1 & c & d \\ 1 & a & b \end{matrix} \right\}_q \end{aligned} \quad (\text{A.4})$$

where

$$\begin{aligned} S_0(\vartheta) &= \pm \frac{1}{4i} \left(\sinh \frac{\pi}{\xi} (\vartheta - \pi i) \sinh \frac{\pi}{\xi} \left(\vartheta - \frac{2\pi i}{3} \right) \right)^{-1} \times \\ &\quad \exp \left(-2i \int_0^\infty \frac{\sin k\vartheta \sinh \frac{\pi k}{3} \cosh \left(\frac{\pi}{6} - \frac{\xi}{2} \right) k}{k \cosh \frac{\pi k}{2} \sinh \frac{\xi k}{2}} dk \right). \end{aligned} \quad (\text{A.5})$$

$$C_a = a(a + 1) \quad (\text{A.6})$$

and the q -6j symbols take the form

$$\begin{aligned} \left\{ \begin{matrix} 1 & a-2 & a-1 \\ 1 & a & a-1 \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a+2 & a+1 \\ 1 & a & a+1 \end{matrix} \right\}_q = 1 \\ \left\{ \begin{matrix} 1 & a+1 & a+1 \\ 1 & a & a \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a & a \\ 1 & a+1 & a+1 \end{matrix} \right\}_q = \frac{\sqrt{[2a+4][2a]}}{[2a+2]} \\ \left\{ \begin{matrix} 1 & a & a+1 \\ 1 & a & a+1 \end{matrix} \right\}_q &= \frac{[2]}{[2a+1][2a+2]} \\ \left\{ \begin{matrix} 1 & a & a-1 \\ 1 & a & a-1 \end{matrix} \right\}_q &= \frac{[2]}{[2a][2a+1]} \\ \left\{ \begin{matrix} 1 & a+1 & a \\ 1 & a & a+1 \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a & a+1 \\ 1 & a+1 & a \end{matrix} \right\}_q = \frac{\sqrt{[2a+4][2a]}}{[2a+2]} \\ \left\{ \begin{matrix} 1 & a & a-1 \\ 1 & a & a+1 \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a & a+1 \\ 1 & a & a-1 \end{matrix} \right\}_q = \frac{\sqrt{[2a-1][2a+3]}}{[2a+1]} \\ \left\{ \begin{matrix} 1 & a & a \\ 1 & a & a+1 \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a & a+1 \\ 1 & a & a \end{matrix} \right\}_q = -\frac{[2]}{[2a+2]} \sqrt{\frac{[2a+3]}{[2a+1]}} \\ \left\{ \begin{matrix} 1 & a & a \\ 1 & a+1 & a \end{matrix} \right\}_q &= \left\{ \begin{matrix} 1 & a+1 & a \\ 1 & a & a \end{matrix} \right\}_q = \frac{[2]}{[2a+2]} \end{aligned}$$

$$\begin{aligned}
\left\{ \begin{array}{ccc} 1 & a & a \\ 1 & a & a-1 \end{array} \right\}_q &= \left\{ \begin{array}{ccc} 1 & a & a-1 \\ 1 & a & a \end{array} \right\}_q = \frac{[2]}{[2a]} \sqrt{\frac{[2a-1]}{[2a+1]}} \\
\left\{ \begin{array}{ccc} 1 & a & a \\ 1 & a-1 & a \end{array} \right\}_q &= \left\{ \begin{array}{ccc} 1 & a-1 & a \\ 1 & a & a \end{array} \right\}_q = -\frac{[2]}{[2a]} \\
\left\{ \begin{array}{ccc} 1 & a & a \\ 1 & a & a \end{array} \right\}_q &= \frac{[2a-1][2a+3]-1}{[2a][2a+2]}
\end{aligned} \tag{A.7}$$

We remark that the square roots in (A.7) carry a sign ambiguity which must be resolved by an appropriate choice of the branches of the square root function. One must consider the square roots of each q -numbers separately and fix a sign for the expressions

$$\sqrt{[n]}, \quad n = 1, 2, \dots \tag{A.8}$$

and then use the selected representative consistently in all formulas. Such a choice of branch is necessary in order for the amplitudes to satisfy the Yang-Baxter equation, R -matrix unitarity and appropriate crossing relations.

B Scattering amplitudes for $\mathcal{M}_{p,p'} + \Phi_{1,5}$

In this appendix we present a corrected version of the S-matrix of $\Phi_{1,5}$ perturbations written down in the paper [4] by one of the authors. The original formulas have misprints and some of them have the wrong normalisation factors. The correct ones can be obtained by imposing crossing symmetry and RU on the amplitudes. The choice of the normalisation factors amount to defining the scalar product of the multi-kink states correctly (i.e. satisfying the constraints imposed by the quantum group symmetry).

In $\Phi_{1,5}$ perturbations, the allowed vacuum sequences are composed of highest weights of the group $U_{q^4}(sl(2))$ where

$$q = \exp\left(\frac{i\pi p'}{4p}\right). \tag{B.1}$$

They are labelled by $j_k = 0, 1/2, \dots, j_{max}$ with $j_{max} = (p-2)/2$. The adjacency rules for a sequence $\{j_1, \dots, j_n\}$ are $|j_{k+1} - j_k| = 0$ or $1/2$ (Figure 7). The zero difference corresponds to a neutral kink, while the nonzero to charged ones. Let us introduce the notation

$$\begin{aligned}
y &= \exp\left(\frac{\pi}{\xi} \vartheta\right), \quad \xi = \frac{4}{3} \frac{\pi p}{p' - 2p}, \\
[z]_4 &= \frac{q^{4z} - q^{-4z}}{q^4 - q^{-4}}.
\end{aligned} \tag{B.2}$$

The scattering amplitudes for charged kinks are (with a sign corrected with respect to [4])

$$S_{ab}^{cd} = \left(- \left(\frac{y^2}{q} - \frac{q}{y^2} - \frac{1}{q} + q \right) \delta_{ac} \left(\frac{[2b+1]_4 [2d+1]_4}{[2a+1]_4 [2c+1]_4} \right)^{1/2} \right)$$




Figure 7: Adjacency rules for kinks in Φ_{15} perturbations

$$+ \left(\frac{y^2}{q^5} - \frac{q^5}{y^2} - \frac{1}{q} + q \right) \delta_{bd} S_0(\vartheta) , \quad (\text{B.3})$$

where $S_0(\vartheta)$ is the function defined in (A.5). For the sign ambiguities occurring as a result of square roots of q -numbers see the remark at the end of Appendix A.

The remaining amplitudes include neutral kinks. The neutral kink-neutral kink scattering, neutral kink-charged kink forward scattering and neutral kink-charged kink reflection were correct in the original paper [4] and are the following

$$\begin{aligned} S_{aa}^{aa} &= \frac{q^6 y^2 + y^2 q^8 - q^8 - q^4 y^2 + y^2 - q^{10} y^2 + y^4 q^2 - y^2 q^2}{y^2 q^5} S_0(\vartheta) , \\ S_{ab}^{ba} &= \frac{(y^2 + q^6)(y^2 - 1)}{y^2 q^3} S_0(\vartheta) , \\ S_{aa}^{ba} &= -\frac{(q^4 - 1)(y^2 + q^6)}{y q^5} S_0(\vartheta) . \end{aligned} \quad (\text{B.4})$$

The amplitudes describing two charged kinks turning into two neutral ones were incorrectly normalised (they include a nontrivial Clebsh-Gordan in the unrestricted theory [4]). The correct form is

$$S_{ab}^{aa} = i \frac{c_{aa}}{c_{ba}} \frac{(q^4 - 1)(y^2 - 1)}{q^2 y} \quad (\text{B.5})$$

For the reverse process the amplitude is

$$S_{aa}^{ab} = i \frac{c_{ab}}{c_{aa}} \frac{(q^4 - 1)(y^2 - 1)}{q^2 y} \quad (\text{B.6})$$

where the normalisation factors c_{ab} are

$$c_{ab} = \begin{cases} \alpha_1 (-1)^{2a} \sqrt{\frac{[2b+1]_4}{[2a+1]_4}} & a = b \pm \frac{1}{2} \\ \alpha_2 & a = b \end{cases} . \quad (\text{B.7})$$

$\alpha_{1,2}$ are constants which are left free by the constraints of RU and crossing. The c_{ab} appear in the crossing relations, which take the form

$$S_{ab}^{cd}(\vartheta) = \frac{c_{ad}}{c_{bc}} S_{da}^{bc}(i\pi - \vartheta) . \quad (\text{B.8})$$

C Transfer matrices and the Bethe-Yang equations

C.1 RSOS transfer matrices

When we put the theory on a cylindrical space-time with spatial volume L , the allowed sequences of vacua (see appendices A and B for ϕ_{12} and ϕ_{15} perturbations, respectively, and [11] for the ϕ_{13} case) are further restricted by the condition $a_1 = a_{N+1}$, where N is the number of particles. We take all particles to be point-like and ignore all vacuum polarisation contributions. Let us define the following (generalised) transfer matrix

$$T_{a_1 a_2 \dots a_N}^{b_1 b_2 \dots b_N}(\vartheta | \vartheta_1, \vartheta_2, \dots, \vartheta_N) = \frac{b_1}{a_1} \left| \begin{array}{c|c} b_2 & \\ \hline a_2 & \end{array} \right| \dots \left| \begin{array}{c|c} b_N & b_1 \\ \hline a_N & a_1 \end{array} \right| \vartheta$$

$\vartheta_1 \quad \vartheta_2 \qquad \qquad \vartheta_N$

which translates to

$$T(\vartheta | \vartheta_1, \vartheta_2, \dots, \vartheta_N)_{a_1 a_2 \dots a_N}^{b_1 b_2 \dots b_N} = \prod_{j=1}^N S_{b_j a_j}^{a_{j+1} b_{j+1}}(\vartheta - \vartheta_j) \quad (\text{C.1})$$

with the identification $a_{N+1} \equiv a_1$, $b_{N+1} \equiv b_1$. In the following we shall not always write down the matrix indices explicitly. From the Yang-Baxter equation, it is straightforward to prove that these transfer matrices form a commuting family

$$[T(\vartheta | \vartheta_1, \dots, \vartheta_N), T(\vartheta' | \vartheta_1, \dots, \vartheta_N)] = 0 \quad (\text{C.2})$$

We define the following specialised transfer matrices

$$\begin{aligned} T_k(\vartheta_1, \vartheta_2, \dots, \vartheta_N)_{a_1 a_2 \dots a_N}^{b_1 b_2 \dots b_N} &= T(\vartheta_k | \vartheta_1, \vartheta_2, \dots, \vartheta_N)_{a_1 a_2 \dots a_N}^{b_1 b_2 \dots b_N} \\ &= (-1)^\delta \delta_{a_k}^{b_{k+1}} \prod_{j \neq k} S_{b_j a_j}^{a_{j+1} b_{j+1}}(\vartheta_k - \vartheta_j) \end{aligned} \quad (\text{C.3})$$

Apart from a phase coming from a plane wave factor of rapidity ϑ_k in the wave function, this gives the monodromy corresponding to taking the k th kink around the spatial circle multiplied by a factor $(-1)^\delta$. The total phase must equal $(-1)^F$ depending on the statistics of the particle k ($F = 1$ for fermions $F = 0$ for bosons). Thus we obtain the so-called Bethe-Yang equations [18]:

$$\exp(im_k R \sinh \vartheta_k) T_k(\vartheta_1, \vartheta_2, \dots, \vartheta_N)_{a_1 a_2 \dots a_N}^{b_1 b_2 \dots b_N} \Psi^{a_1 a_2 \dots a_N} = (-1)^{F+\delta} \Psi^{b_1 b_2 \dots b_N} \quad (\text{C.4})$$

where $\Psi^{a_1 a_2 \dots a_N}$ is the wave function amplitude, defined by the decomposition of the state $|\Psi\rangle$ as follows

$$|\Psi\rangle = \sum_{a_1, \dots, a_N} \Psi^{a_1 a_2 \dots a_N} |K_{a_1 a_2}(\vartheta_1) \dots K_{a_N a_1}(\vartheta_N)\rangle. \quad (\text{C.5})$$

The energy and the momentum of the state (relative to the vacuum) are given by

$$E = \sum_{k=1}^N m_k \cosh \vartheta_k, \quad P = \sum_{k=1}^N m_k \sinh \vartheta_k. \quad (\text{C.6})$$

Due to the commutation relation (C.2), the equations (C.4) for the vector Ψ are compatible and can be reduced to scalar equations by simultaneously diagonalising the transfer matrices. Let us denote the eigenvalues of $T(\vartheta|\vartheta_1, \dots, \vartheta_N)$ by $\lambda^{(s)}(\vartheta|\vartheta_1, \dots, \vartheta_N)$ with the corresponding eigenvectors $\psi^{(s)}(\vartheta_1, \dots, \vartheta_N)$ (s is just an index enumerating the eigenvalues and the eigenvectors can be chosen independent of ϑ due to the commutativity (C.2)). Then the solutions of the Bethe-Yang equations (C.4) are given by

$$\Psi^{a_1 a_2 \dots a_N} = \psi^{(s)}(\vartheta_1, \dots, \vartheta_N)^{a_1 a_2 \dots a_N} \quad (\text{C.7})$$

where the rapidities solve the scalar Bethe Ansatz equations

$$\exp(im_k R \sinh \vartheta_k) \lambda^{(s)}(\vartheta_k|\vartheta_1, \dots, \vartheta_N) = (-1)^{F+\delta} \quad (\text{C.8})$$

C.2 Folded transfer matrices

For folded models the vacua are labelled by an integer a modulo the folding number k . The allowed sequences satisfy

$$a_{i+1} = a_i + Q \pmod k \quad (\text{C.9})$$

where Q are the possible topological charges of the solitons

$$Q = \begin{cases} \pm 1, & \text{sine - Gordon} \\ +1, 0, -1 & \text{ZMS} \end{cases} \quad (\text{C.10})$$

In finite volume we require $a_1 = a_{N+1} \pmod k$ for periodic boundary conditions. Then the transfer matrix has the same form as in equation (C.1) except that now the matrix S is constructed from the scattering matrix $\tilde{S}$ of the sine-Gordon/ZMS model in the following way:

$$S_{ab}^{cd}(\vartheta) = \tilde{S}_{Q_{ab}, Q_{bc}}^{Q_{ad}, Q_{dc}}(\vartheta) \quad (\text{C.11})$$

where Q_{ab} denotes the charge of the soliton connecting the vacua a and b (see Figure 8).




Figure 8: Vacuum labels and topological charges for two-particle S -matrices in folded models

References

- [1] C. Mudry, C. Chamon and X.-G. Wen, *Nucl. Phys.* **B 466** (1996) 383-443, cond-mat/9509054.
N. Read and H. Saleur, *Nucl. Phys.* **B 613** (2001) 409-444, hep-th/0106124.
M. J. Bhaseen, J.-S. Caux, I. I. Kogan and A. M. Tsvelik, *Nucl. Phys.* **B 618** (2001) 465-499, cond-mat/0012240.
- [2] G. Takács and G. Watts, *Nucl. Phys.* **B547** (1999) 538-568, hep-th/9810006.
- [3] P. Dorey and R. Tateo, *Nucl. Phys.* **B 571** (2000) 583-606, hep-th/9910102.
- [4] G. Takács, *Nucl. Phys.* **B 489** (1997) 532-556, hep-th/9604098.
- [5] J.L. Cardy and G. Mussardo, *Phys. Lett.* **B225** (1989) 275-278.
- [6] H.G.Kausch, G.Takács and G.M.T.Watts, *Nucl. Phys.* **B 489** (1997) 557-579, hep-th/9605104.
- [7] D.I. Olive, *Nuovo Cimento* **26** (1962) 73.
- [8] J.L. Miramontes, *Phys. Lett.* **B455** (1999) 231-238, hep-th/9901145.
- [9] T.R. Klassen and E. Melzer, *Nucl. Phys.* **B 338** (1990) 485-528.
- [10] Z. Bajnok, L. Palla, G. Takács and F. Wágner, *Nucl. Phys.* **B 587** (2000) 585-618, hep-th/0004181.
- [11] N. Reshetikhin and F.A. Smirnov, *Commun. Math. Phys.* **131** (1990) 157-178.
- [12] F.A. Smirnov, *Int. J. Mod. Phys.* **A6** (1991) 1407-1428.
- [13] A. Klümper and P.A. Pearce, *J. Stat. Phys.* **64** (1991) 13;
A. Klümper, M. Batchelor and P.A. Pearce, *J. Phys.* **A24** (1991) 3111.
- [14] C. Destri and H.J. de Vega, *Phys. Rev. Lett.* **69** (1992) 2313; *Nucl. Phys.* **B 438** (1995) 413-454, hep-th/9407117.

- [15] D. Fioravanti, A. Mariottini, E. Quattrini and F. Ravanini, *Phys. Lett.* **B 390** (1997) 243-251, hep-th/9608091.
- [16] C. Destri and H.J. de Vega, *Nucl. Phys.* **B 504** (1997) 621-664, hep-th/9701107.
- [17] G. Feverati, F. Ravanini and G. Takács, *Phys. Lett.* **B430** (1998) 264-273, hep-th/9803104; *Nucl. Phys.* **B 540** (1999) 543-586, hep-th/9805117; *Phys. Lett.* **B444** (1998) 442-450, hep-th/9807160.
- [18] T.R. Klassen and E. Melzer, *Nucl. Phys.* **B 382** (1992) 441-485, hep-th/9202034.
- [19] P.E. Dorey and R. Tateo, *Nucl. Phys.* **B 482** (1996) 639-659, hep-th/9706140; *Nucl. Phys.* **B 515** (1998) 575-623, hep-th/9607167.
- [20] Al. B. Zamolodchikov, *Nucl. Phys.* **B 432** (1994) 427-456; *Phys. Lett.* **B335** (1994) 436-443.
- [21] G. Feverati, F. Ravanini and G. Takács, *Nucl. Phys.* **B 570** (2000) 615-643, hep-th/9909031.
- [22] P. Zinn-Justin, *J. Phys.* **A31** (1998) 6747-6770, hep-th/9712222
- [23] S. Coleman and H.J. Thun *Commun. Math. Phys.* **61** (1978) 31.
- [24] V.B. Lidskii, *U.S.S.R. Comput. Math. and Math. Phys.* **6** (1966) 73-85, J. Moro, J.V. Burke and M.L. Overton, *SIAM Journal on Matrix Analysis and Applications* **18** (1997) 793-817.
- [25] G. Mussardo, *Int. J. Mod. Phys.* **A7** (1992) 5027-5044.
- [26] F. Ravanini, M. Stanishkov and R. Tateo *Int. J. Mod. Phys* **A11** (1996) 677-698, hep-th/9411085.
- [27] M.J. Martins, *Phys. Rev. Lett.* **69** (1992) 2461-2464, hep-th/9205024; *Nucl. Phys. B* **394** (1993) 339-355.
- [28] E. Melzer, *Supersymmetric Analogs of the Gordon-Andrews Identities, and Related TBA Systems*, hep-th/9412154.
- [29] P.E. Dorey, C. Dunning and R. Tateo, *Nucl. Phys.* **B 578** (2000) 699-727, hep-th/0001185.
- [30] P.E. Dorey, A. Pocklington and R. Tateo, Integrable aspects of the scaling q -state Potts models I: bound states and bootstrap closure *and* Integrable aspects of the scaling q -state Potts models II: finite size effects (to appear).